

## Delayed surgical intervention:

In most cases swelling and bruising will start to lessen after a week but fractured bones will take longer to heal. Swelling can make it difficult to examine the eye and its movement – your doctor may decide to re-assess you after a week when the swelling has reduced.

If delayed surgery is decided upon it will be for the following reasons:

- To remove bony fragments
- To get rid of double vision and free trapped tissue
- To restore the normal shape of the eye socket and eye position

Surgery is always a compromise between the best access to the fracture and the potential complications of each approach. Exactly what surgery is done depends on the position of the defect, the age of the person and the existence of other related injuries. Your surgeon will discuss what surgical approach is best if you need to have surgery. If double vision persists after surgery another Orthoptic assessment will need to be done – it is possible to help reduce double vision using prisms and/or a patch.

## Contact numbers

### Orthoptist:

**St Richard's**  
**01243 788122 ext 3514**

**Worthing**  
**01903 205111 ext 85430**

**We are committed to making our publications as accessible as possible. If you need this document in an alternative format, for example, large print, Braille or a language other than English, please contact the Communications Office by:  
email: [Communications@wsht.nhs.uk](mailto:Communications@wsht.nhs.uk)  
or by calling 01903 205 111 ext 84038.**

[www.westernsussexhospitals.nhs.uk](http://www.westernsussexhospitals.nhs.uk)

Department: Orthoptics  
Issue date: **Nov 2013**  
Review date: **Nov 2015**  
Leaflet Ref: ORT19



## Orthoptic Department Information Sheet

### Orbital Fractures

St Richard's  
Hospital  
Spitalfield Lane  
Chichester  
West Sussex  
PO19 6SE

Worthing Hospital  
Lyndhurst Road  
Worthing  
West Sussex  
BN11 2DH

This leaflet is intended to answer some of the questions of patients or carers of patients, diagnosed with orbital fracture under the care of Western Sussex Hospitals NHS Trust.

### **What is an orbital fracture?**

The eye socket, or orbit, is a bony cup that surrounds and protects the eye. When one of the bones that forms this bony cup is broken it is called an orbital fracture.

### **How is it caused?**

Road traffic accidents, assault, sports injuries, industrial injuries and falls are all common causes of this type of facial fracture. Some injuries of this nature are preventable by the use of seatbelts when driving and using protective eye wear when at work or playing some sports.

### **What are the signs I have a fracture?**

Signs and symptoms are going to vary according to the cause and severity of the injury. Commonly experienced problems are:

Pain - to the eye and the soft tissue around it.

Double vision - swelling, bleeding or entrapment of the soft tissue or muscles around the eye can restrict its movement. A nerve injury can also affect the eye's movement. Sometimes the eyes can turn or 'squint'. When the eyes are unable to move as a synchronised pair this can lead to double vision.

Blurred vision - a direct blow to the eyeball can cause damage affecting the vision. Depending on the nature of the injury this could be mild and heal itself such as an abrasion to the cornea (front clear surface of the eye), or be severe enough to cause a cataract or traumatic damage to the nerve of the eye.

Loss of sensation - a fracture of the bony wall of the eye socket can result in damage to the nerve responsible for sensation to the cheek, eyelids, upper lip and teeth or the forehead.

Bruising - a black eye with bruising around the injured area. There can be areas of redness on the white of the eye and on the inner lining of the eyelids.

Puffy eyelids - a puffy large amount of air under the skin around the eye can be a sign that the bony wall between the eye and the sinus is broken. Do not blow your nose while the eye heals to lessen the chance of this air accumulation from happening.

Sunken eye - a broken bone in the eye socket can cause some of the soft tissue in the socket to leak into other areas such as the sinuses. This loss of 'packing' tissue can cause the eye to look like it has sunken or dropped.

### **How is it tested?**

Testing will usually involve visiting more than one department.

The Maxillo-facial Department - specialises in the diagnosis and treatment of diseases

and injuries to the mouth, face and neck. This department co-ordinates testing and management (including surgery if needed). Radiology Department - a CT or MRI scan enables the doctors to see the extent of bony and soft tissue injury.

Eye Clinic - if it is suspected that your vision is reduced or that you have a wound or infection to your eyeball or eyelids then an Ophthalmologist will be asked to do an examination and help with the management of the eye health.

Orthoptic Department - if double vision is noticed then an Orthoptist will be asked to do a map of the eye movements (a Hess chart) and measure both the eye position and how well they work together.

### **How can it be treated?**

Treatment depends on the severity and location of the injury.

Conservative treatment: For a small uncomplicated fracture with minimal double vision the doctor may advise pain relief, cold compression to reduce the swelling and a few days rest. Do not blow the nose whilst the eye heals.

Emergency surgical intervention: If the fracture is more severe, surgical repair might need to be done - the speed of which depending on how bad the injury is. Entrapped muscle can lead to nausea, vomiting and a low heart rate, especially in children. These symptoms will result in urgent surgery - usually within 4 days.